


Form of emergency contraception	Chance of pregnancy (will be affected by place in menstrual cycle)	How soon do I need to use it after unprotected sex?	How do I use it?	Extra info
 IUD (copper coil)	 Most effective- almost 100%.	 <b>5</b> Up to 5 days (120 hours-longer in certain cases)	Inserted into the uterus (womb) by a health professional in a clinic	 Provides ongoing contraception for up to 10 years
Ullipristal acetate (ellaone)	 2nd most effective	 Up to 5 days (120 hours)	 A tablet	Unlikely to work if you have already released an egg from your ovary (ovulated)
Levonergestrel (plan B)	 Least effective	 Up to 3 days (72 hours)	 A tablet	Unlikely to work if you have already released an egg from your ovary (ovulated)

Please note, exceptions apply- every patient and situation is different

# Effectiveness of Emergency Contraception

1000 women have a single episode of UPSI



If EC not used  
55 of 1000  
become pregnant

If all 1000 use  
Levonelle 22  
become pregnant

If all 1000 use  
EllaOne14 become  
pregnant

If all 1000 have  
emergency IUD  
1 becomes pregnant

OR 0.58; 95% CI 0.33–0.99,  $p=0.046$ .  
1.4% (22/1617 pregnancies), 2.2% (35/1625 pregnancies).

Glazier AF et al., *Lancet* 2010; **375**: 555–62  
Cleland K et al *Human Reprod* 2012

Rachel D'Souza