



Central and  
North West London  
NHS Foundation Trust

## Important information about your medication

You have been prescribed a medication that interacts with some common supplements and medications. Taking these tablets or liquids can result in your medication not working as effectively. Some of these are prescribed and others can be purchased over the counter from a pharmacy.

# Rilpivirine (Edurant, Eviplera, Juluca and Odefsey)

Proton pump inhibitors, for example, Omeprazole

Cannot be taken together

H2 antagonists, for example, Ranitidine

Take 12 hours **before** or at least four hours **after** Rilpivirine

Antacids, for example, Rennie and Gaviscon

Take antacids at least two hours **before** or four hours **after** Rilpivirine

Multivitamins

Calcium supplements

No interaction – can be taken together

## Atazanavir (Reyataz and Evotaz)

Proton pump inhibitors, for example, Omeprazole

Should not be taken together

H2 antagonists, for example, Ranitidine

Caution required – discuss with pharmacy

Antacids, for example, Rennie and Gaviscon

Take Atazanavir at least two hours **before** or one hour **after** antacids

The medications listed can be bought under many different names. Please always check the label or ask the pharmacist. **If in doubt, check with us before starting new medications.**

There may be interactions with other medications, please visit **[www.hiv-druginteractions.org](http://www.hiv-druginteractions.org)** for more information

Tel: 020 3317 5228

[www.cnwl.nhs.uk](http://www.cnwl.nhs.uk)